

Routine Immunization Schedules for Children, Youth & Adults

The vaccines listed on these schedules are given free of charge.

Children, youth and adults with high-risk conditions may require additional doses and be eligible for additional vaccines. For more information, talk to your health care provider or call your Public Health office.

| Childhood Vaccines | | Schedule | | | | | |
|---|----------|----------|-----------------|-----------|-----------|------------------|--|
| | 2 months | 4 months | 6 months | 12 months | 18 months | 4-6 years | |
| Influenza* | | | * | | | | |
| DTaP-IPV-Hib Diphtheria, tetanus, acellular pertussis (whooping cough), polio, and Haemophilus influenzae type b vaccine | / | ~ | ~ | | / | | |
| Pneumo Conj. Pneumococcal conjugate vaccine | V | / | | / | | | |
| Men C Conj. Meningococcal group C conjugate vaccine | | | | / | | | |
| MMRV** Measles, mumps, rubella and varicella vaccine | | | | / | ** | | |
| Tdap-IPV Tetanus, diphtheria, acellular pertussis (whooping cough), and polio vaccine | | | | | | / | |

^{*} Every flu season for all children 6 months of age and older. Children under 9 years old getting their first flu vaccine need 2 doses.

^{**} The second dose of MMRV can be given between 18 months and 6 years of age.

| School Vaccines | Grade 7 |
|--|----------|
| HPV Human papillomavirus vaccine | ✓ |
| Hepatitis B Hepatitis B vaccine | ✓ |
| Tdap Tetanus, diphtheria, and acellular pertussis (whooping cough) vaccine | ✓ |
| Meningococcal Quadrivalent Meningococcal Quadrivalent vaccine (Groups A, C, Y & W 135) | ✓ |

| A -ll. \ | Schedule | | | |
|---|------------|-------------------|--|--|
| Adult Vaccines | All Adults | Adults 65 & older | | |
| Influenza Every flu season | ✓ | | | |
| High-dose Influenza* | | * | | |
| Td Tetanus, diphtheria vaccine (Every 10 years) | ✓ | | | |
| Tdap** Tetanus, diphtheria and acellular pertussis vaccine (If not previously given in adulthood, one dose 10 years after receipt of a tetanus containing vaccine.) | ✓ | | | |
| Pneumococcal polysaccharide Pneumococcal polysaccharide vaccine | | ✓ | | |
| MMR Measles, mumps and rubella vaccine (2 doses for those born in 1970 or later) | ✓ | | | |

^{*} Every flu season for Long-Term Care Facility (Nursing Home and Residential Care Facility) residents only.

^{**} Regardless of age, Tdap should be offered with every pregnancy.